

In silico identification of novel selenoproteins in the *Drosophila melanogaster* genome

Sergi Castellano, Nadya Morozova¹, Marta Morey², Marla J. Berry¹, Florenci Serras², Montserrat Corominas² & Roderic Guigó⁺

Grup de Recerca en Informàtica Biomèdica, Institut Municipal d'Investigació Mèdica (IMIM), Universitat Pompeu Fabra, Dr. Aiguader 80, 08003 Barcelona, ²Departament de Genètica, Universitat de Barcelona, Diagonal 645, 08071 Barcelona, Spain and ¹Thyroid Division, Harvard Institutes of Medicine, 77 Avenue Louis Pasteur, Boston, MA 02115, USA

Received April 4, 2001; revised May 31, 2001; accepted June 6, 2001

In selenoproteins, incorporation of the amino acid selenocysteine is specified by the UGA codon, usually a stop signal. The alternative decoding of UGA is conferred by an mRNA structure, the SECIS element, located in the 3'-untranslated region of the selenoprotein mRNA. Because of the non-standard use of the UGA codon, current computational gene prediction methods are unable to identify selenoproteins in the sequence of the eukaryotic genomes. Here we describe a method to predict selenoproteins in genomic sequences, which relies on the prediction of SECIS elements in coordination with the prediction of genes in which the strong codon bias characteristic of protein coding regions extends beyond a TGA codon interrupting the open reading frame. We applied the method to the *Drosophila melanogaster* genome, and predicted four potential selenoprotein genes. One of them belongs to a known family of selenoproteins, and we have tested experimentally two other predictions with positive results. Finally, we have characterized the expression pattern of these two novel selenoprotein genes.

INTRODUCTION

Selenoproteins are proteins that incorporate the amino acid selenocysteine, a cysteine analog in which a selenium atom is found in place of sulfur. Several components of the selenoprotein synthesis machinery are conserved between different species, suggesting an important role of selenoproteins in cell function (Low and Berry, 1996; Stadman, 1996). Incorporation of selenocysteine into selenoproteins requires an unusual translation step where UGA, normally a stop codon, specifies selenocysteine insertion. Thus, in a single mRNA, UGA can have two contrasting meanings: stop or selenocysteine. The

alternative decoding of UGA is conferred by an mRNA secondary/tertiary structure (the selenocysteine insertion sequence, the SECIS element), which is located in eukaryotes in the 3'-untranslated region. SECIS structures are divided into two classes, termed form 1 and form 2, the latter having an additional small stem-loop at the top of the SECIS element. Most selenoproteins contain a single selenocysteine residue per polypeptide chain, but selenoprotein P has as many as 10–12 (Tujebajeva *et al.*, 2000a).

Selenoproteins have been identified in Bacteria, Archaea and Eukarya. Among eukaryotes, selenoproteins appear to be more common in mammals. Thus, 19 selenoproteins have been found to date in mammals (Flohé *et al.*, 2000), but none in the genome of *Saccharomyces cerevisiae*, and only one in the genome of *Caenorhabditis elegans* (Buettner *et al.*, 1999; Gladyshev *et al.*, 1999). Recently, the class 2 selenophosphate synthetase gene (*sps2*)—an enzyme in the pathway of selenoprotein synthesis, and a selenoprotein itself in mammals—has also been shown to be a selenoprotein in *Drosophila melanogaster* (Hirosawa-Takamori *et al.*, 2000). So far, it remains the only selenoprotein identified in this organism and maps to chromosome 2L. However, pupal proteins of 68, 42 and 25 kDa have been reported to incorporate selenium (Robinson and Cooley, 1997), and a major band of 42 kDa has also been observed in protein extracts of larvae labeled with ⁷⁵Se (Alsina *et al.*, 1999). In addition, some components of the selenoprotein synthesis machinery have already been identified in the fly (Persson *et al.*, 1997; Alsina *et al.*, 1998; Zhou *et al.*, 1999). Moreover, a mutation in the *sps1* gene (a cysteine homolog of *Sps2*) leads to larval lethality, increased apoptosis and aberrant imaginal disc morphology (Alsina *et al.*, 1998). These data strongly suggest the existence of as yet

*Corresponding author. Tel: +34 93 225 7567; Fax: +34 93 221 3237; E-mail: rguigo@imim.es

S. Castellano *et al.*

unidentified selenoproteins in the *D. melanogaster* genome. The recent availability of the complete DNA sequence of this genome should constitute an invaluable resource for characterizing the *D. melanogaster* selenoproteins.

Prediction of selenoproteins in genomic sequences, however, is particularly difficult. Without exception, computational gene prediction programs rely on the standard stop codons TAA, TAG and TGA to identify open reading frames (ORFs) and predict coding exons, through the determination of suitable splicing sites and the computation of some measure of coding likelihood, usually related to bias in codon usage (see Burge and Karlin, 1998 and Haussler, 1998 for reviews on computational gene finding). Under such an assumption, selenoprotein genes, in which TGA does not necessarily imply termination of translation, will be incorrectly predicted. Indeed, the *D. melanogaster sps2* gene (*dsps2*) is wrongly predicted in the released annotation of the fly genome: >100 amino acids are missing from a 379 amino acids protein. Correct delineation of the exonic structure is singularly important to predict selenoprotein genes. Misprediction of only a single amino acid (the selenocysteine residue) may lead to misidentification of selenoproteins (see Results).

Although searching for potential SECIS elements has proved useful in identifying new selenoproteins in expressed sequence tag (EST) sequences (Kryukov *et al.*, 1999; Lescure *et al.*, 1999), this approach is impractical when applied to genomic sequences, given the high frequency of occurrence of the SECIS pattern (see Results). To reduce the number of false positive predictions, we developed a method that relies on the correlated prediction of SECIS elements and of genes in which the strong codon bias characteristic of protein coding regions extends beyond a TGA codon interrupting the ORF. Indeed, we have found that in selenoproteins the region comprised between the in-frame TGA codon and the stop codon shows codon bias comparable to that found in coding regions, while in non-selenoproteins the region comprised between the stop codon TGA and the next stop codon in-frame shows codon bias comparable to that in non-coding regions (Supplementary data). Therefore, measures of codon bias can be used to distinguish actual selenoproteins from false predictions in SECIS-positive nucleotide sequences

RESULTS

Prediction of novel selenoproteins in the *D. melanogaster* genome

The March 24, 2000 release of the *D. melanogaster* genome sequence summing up 115 229 998 bp and containing 13 329 annotated genes was used (Adams *et al.*, 2000). 37 876 potential SECIS elements were found along this sequence using the program PatScan. The minimum free energy of each putative SECIS was measured, and only those fitting an energy stability criteria were considered further. This resulted in 1220 potential SECIS. Along with the sequence, positions of these elements were given to a modified version of the program geneid, which allows for the prediction of genes interrupted by in-frame TGA codons. The restriction was enforced such that genes could not be further than 500 bp upstream from a predicted SECIS. Eleven potential selenoproteins were predicted among a total of 12 194 genes. Seven of them were discarded because the predicted

exonic structure was incompatible with the exonic structure of known overlapping genes, contradicted identical EST sequences or was similar to known proteins with functions apparently unrelated to those of selenoproteins. Of the remaining four, one corresponded to the previously identified *D. melanogaster* selenoprotein *dsps2*. For another two, we identified cysteine paralogs within the set of proteins predicted in the *D. melanogaster* genome. No additional evidence was found for the fourth putative selenoprotein, after an exhaustive search against a number of databases of known coding sequences using the BLAST suite of programs (Altschul *et al.*, 1997). In addition, the predicted secondary structure around the selenocysteine residue of this putative selenoprotein is not compatible with the known crystal structure of the bovine glutathione peroxidase, a eukaryotic selenoprotein. This structure appears to be common to most known selenoproteins, including the two other predictions (Supplementary data). We have thus considered this prediction to be a false positive. Incorporation of selenium was subsequently demonstrated for the two other predicted selenoproteins (which we name *dseI*G and *dseI*M), and their expression pattern during development was characterized.

*dseI*G

*dseI*G is a 110 aa protein which maps to 10F4-6 of the X chromosome. It differs in only two amino acids from the annotated protein in *D. melanogaster* (CG1844), the in-frame TGA lying only one codon upstream from the stop codon. *dseI*G has a cysteine paralog, the *CG1840* gene. They appear in tandem, separated by only 320 bp, and have the same exonic structure sharing 65% identity at the protein level (Figure 1).

*dseI*M

*dseI*M is a 249 aa protein which maps to 12A4-6 of the X chromosome. It differs substantially from the protein annotated in *D. melanogaster* (CG11177), the first exon, and a large fraction of the second (in which the in-frame TGA is located) having been missed. *dseI*M has two distant paralogs, the *CG13186* and *CG15147* genes (Figure 2).

⁷⁵Se labeling of the *D. melanogaster* selenoproteins expressed in mammalian cells

⁷⁵Se labeling of HEK cells was undertaken to demonstrate that *GH03581* (*dseI*G) and *SD09114* (*dseI*M) genes encoded bona fide selenoproteins. In cells transfected with empty vector, the background pattern of endogenously expressed selenoproteins can be seen (Figure 3A, lane 1), including thioredoxin reductases (~55 kDa), glutathione peroxidase (~23 kDa), phospholipid hydroperoxide glutathione peroxidase (~20 kDa), and an ~12–14 kDa triplet. Transfection of the *GH03581* expression vector resulted in an increase in labeling in the 12 kDa size range, overlapping the lower band of the endogenous triplet (Figure 3A, lane 2). Transfection of the *SD09114* expression vector resulted in appearance of a prominent new band of ~30 kDa (Figure 3A, lane 3), corresponding to the predicted size of *SD09114* (~27 kDa). *SD09114* protein appears to be a *D. melanogaster*-specific selenoprotein, with no analog in mammalian cells of similar molecular weight.




Fig. 1. *dSelG*. (A) Gene structure for *dSelG* and in-tandem *CG1840* paralog plotted using gff2ps (Abril and Guigó, 2000). Coordinates correspond to the AE002593 (X) scaffold. The extra coding region is shown in red as predicted by geneid and the annotated coding exons are in blue. (B) *dSelG* form 2 SECIS. (C) Alignment of *dSelG* and *CG1840* paralogs using CLUSTAL_W (Thompson *et al.*, 1994).




Fig. 2. *dSelM*. (A) Gene structure for *dSelM* and single exon *CG15147*, *CG13186* paralogs. Coordinates correspond to the AE002593 (X), AE002690 (2L) and AE002787 (2R) scaffolds, respectively. (B) *dSelM* form 2 SECIS. (C) Alignment of *dSelM* and *CG15147*, *CG13186* paralogs.

In situ hybridization in embryos, discs and brains

In situ hybridization experiments were performed to assess *dSelG* and *dSelM* mRNA expression patterns. *dSelM* mRNA was present in all embryonic stages, especially in the blastoderm stage, suggesting that there is a strong maternal contribution (Figure 4A, C and E). Imaginal discs displayed a ubiquitous *dSelM* expression pattern (Figure 4I), and in brain, although the staining was ubiquitous, large cells, probably neuroblasts, were highly stained (Figure 4G). *dSelG* expression pattern was analyzed in embryos, and similarly to *dSelM*, the mRNA was found ubiquitously in all stages (Figure 4K, M and O). Due to the high

similarity at the nucleotide sequence level between *dSelG* and the cysteine homolog (64% in the coding fraction), we have to assume that the probe used for the *in situ* hybridization would detect both transcripts, if present.

DISCUSSION

Most of the functions of selenium involve its incorporation into selenoproteins in the form of selenocysteine. Besides their putative role in regulating the redox state of the cell, selenoproteins seem to possess anticarcinogenic properties, and PHGPx plays a role in reproductive function (Ganther, 1999; Ursini *et al.*, 1999).

S. Castellano et al.




Fig. 3. ^{75}Se -labeling of the *D. melanogaster* selenoproteins expressed in mammalian cells. (A) Lane 1: ^{75}Se -labeling of cells transfected with empty vector. Lane 2: ^{75}Se -labeling of cells transfected with *dSelG*. Lane 3: ^{75}Se -labeling of cells transfected with *dSelM*. (B) High magnification of the region corresponding to the *dSelG* labeling.




Fig. 4. *In situ* hybridization in embryos, imaginal discs and brain. (A, C and E) *dselM* expression pattern in syncytial blastoderm, cellular blastoderm and gastrulation embryonic stages, respectively; (B, D and F) the corresponding sense controls (scale bar is 50 mm); (G) *dselM* expression in brain and neuroblast staining in the inset; (H) the brain sense control (scale bar 100 mm, inset scale bar 2.5 mm); (I) *dselM* expression in wing disc; (J) wing disc sense control (scale bar 50 mm); (K, M and O) *dselG* expression pattern in syncytial blastoderm, cellular blastoderm and gastrulation embryonic stages, respectively; (L, N and P) are the corresponding sense controls (scale bar is 50 mm).

Drosophila provides a convenient tool for investigating selenoprotein function because of the availability of fly genetics and the already existing mutation in the *sps1* gene (Alsina et al., 1998).

Using a novel computational method we have predicted four potential selenoprotein genes in the *D. melanogaster* genome,




Fig. 5. SECIS and gene prediction. (A) General form 1 SECIS divided into structural units. Form 2 has an extra short stem-loop in the apical loop. (B) PatScan SECIS pattern to search for both form 1 and form 2 SECIS. The extra stem-loop in form 2 is not taken into account when searching. (C) The two possible ways of geneid prediction for an ideal two exons gene: as a normal gene or as a selenoprotein gene with a TGA in-frame and a SECIS. Exon defining signals are shown. (D) False positive selenoprotein genes with either a TGA in-frame or a SECIS. These partial predictions are not permitted in the gene prediction.

with little human intervention. Three are bona fide selenoproteins: *dSps2*, already demonstrated as such (Hirosawa-Takamori et al., 2000), and the other two, *dSelG* and *dSelM*, shown herein by ^{75}Se labeling. In addition, cysteine paralogs exist in *D. melanogaster* for these three selenoproteins. While *Sps2*, a selenophosphate synthetase, belongs to a known family of selenoproteins, *dSelG* and *dSelM* are novel selenoproteins, lacking sequence similarity to known proteins. *dSelG* has a cysteine homolog in *C. elegans* of unknown function, while *dSelM* appears to belong to a new class of selenoproteins widely distributed across the phylogenetic spectrum: we have found selenocysteine homologs to *dSelM* in ESTs from zebrafish, human and mouse databases, among other organisms.

It is unclear, however, how complete our characterization of the selenoprotein set in *D. melanogaster* is. Experimental data suggest the existence of a selenoprotein in the 60–70 kDa range (Alsina et al., 1999), for which we have not been able to find a computational prediction. Forcing the SECIS element to occur within 500 bp downstream from the selenoprotein coding region may be too restrictive. Although most mammalian selenoproteins are within this range, longer distances up to >4000 bp are possible. The fly genome is certainly more compact, and while in the human *sps2* the SECIS element is 579 bp downstream from the stop codon, this distance is only 30 bp in *D. melanogaster*. On the other hand, exceptions to the standard eukaryotic SECIS model have recently been reported in *C. elegans*. In this case, a 5'-GUGA motif is present instead of AUGA (Buettner et al., 1999). Therefore, it is possible that additional selenoproteins using an alternative




Fig. 6. General schema for selenoprotein identification.

SECIS structure exist in the *D. melanogaster* genome. Relaxing the SECIS pattern to capture a more general SECIS structure results in a substantial increase in the number of predicted SECIS elements, which compounds the analysis of the search results. In this regard, the approach presented here could contribute towards systematically exploring alternative SECIS structures.

In summary, we believe that the research described here demonstrates the power of the combined *in silico*, *in vitro* and *in vivo* approaches towards a better understanding of living systems.

METHODS

Prediction of selenoproteins in nucleotide sequences. The method that we have developed is described in detail in the Supplementary data, which can be found at *EMBO reports* Online. A general schema is shown in Figure 6. Broadly, given a query sequence, first we predict SECIS elements using the program PatScan (<http://www-unix.mcs.anl.gov/compbio/PatScan/HTML/PatScan.html>) (Figure 5A and B). The stability of the predicted SECIS is then assessed using the RNAfold program (Vienna RNA package) using the protocol by Kryukov *et al.* (1999). Next, we use a modification of the program geneid (Guigó *et al.*, 1992; Parra *et al.*, 2000) to predict genes that may be interrupted by in-frame TGA codons. Such genes, however, can be predicted only when a putative SECIS, whose position along the genome is input into geneid during gene prediction, exists at the right distance. The modified geneid yields, in the same gene prediction, both standard genes and selenoprotein genes (Figure 5C and D).

⁷⁵Se labeling. The pOT2 plasmids containing GH03581 (*dseI/G*) and SD09114 (*dseI/M*) cDNA clones were obtained from Research Genetics Inc. and sequenced using the Dye Terminator

Cycle Sequencing method. Inserts were subcloned into pUHD10–3 vector via *EcoRI* and *XbaI* sites. Human embryonic kidney cells (HEK-293) were transiently transfected by CaPO₄ DNA precipitation method (Tujebajeva *et al.*, 2000b) with either *dseI/G* or *dseI/M* expression plasmids and co-transfected with plasmids encoding tRNA^{[Ser]Sec} (Lee *et al.*, 1990) and SECIS-binding protein (SBP2) (Copeland *et al.*, 2000) to increase the efficiency of selenocysteine incorporation (Berry *et al.*, 1994; Tujebajeva *et al.*, 2000b). All transfection experiments were carried out with supplementation of 100 nM sodium selenite to the media. ⁷⁵Se as sodium selenite (1000 mCi/mg) was added to media 1 day after transfection, and labeling proceeded for another day. Cells were harvested, sonicated in 0.25 M sucrose in PE buffer (0.1 M potassium phosphate, 1 mM EDTA pH 6.9) and analyzed by polyacrylamide gel electrophoresis, followed by autoradiography.

Whole-mount *in situ* hybridization. Embryos collected from a 24-h egg-lay were dechorionated and fixed in 2% formaldehyde and 0.5 M final concentration of EGTA in PBS for 20 min. After precipitation with methanol embryos were kept at –20°C in absolute ethanol. Third-instar wild-type larvae were dissected in PBS and fixed overnight in 4% paraformaldehyde in PBS for 20 min. Further steps before hybridization and hybridization itself were performed as described by Lehner and O’Farrell (1990). Linearized pOT2 vectors containing GH03581 and SD09114 clones were used to generate a riboprobe according to the Boehringer-Mannheim protocol. Embryos and discs were then incubated with 1/2000 anti-DIG conjugated with alkaline phosphatase antibody (Boehringer-Mannheim), preabsorbed against fixed and dissected larvae. Antibody was detected using standard procedures (Boehringer-Mannheim). Embryos were postfixed and posteriorly mounted in DePeX. Discs and brains were dissected and mounted in 87% glycerol.

Data and software availability. Sequence data and software can be found at <http://www1.imim.es/databases/spdroso2001>

Supplementary data. Supplementary data are available at *EMBO reports* Online.

ACKNOWLEDGEMENTS

We wish to thank E. Blanco for his assistance with geneid, and P. Higgs for helpful suggestions. R.G. thanks L.S. Shashidhara for his hospitality at the CCMB (Hyderabad), where part of this manuscript was written. This work was partially supported by grants BIO98-0443-C02-01 and PB96-1253 from ‘Plan Nacional de I+D’ (Spain). S.C. and M.M. are recipients of predoctoral fellowships from CIRIT, Generalitat de Catalunya.

REFERENCES

- Abril, J.F. and Guigó, R. (2000) gff2ps: visualizing genomic annotations. *Bioinformatics*, **16**, 743–744.
- Adams, M.D. *et al.* (2000) The genome sequence of *Drosophila melanogaster*. *Science*, **287**, 2185–2195.
- Alsina, B., Serras, F., Bagaña, J. and Corominas, M. (1998) Patufet, the gene encoding the *Drosophila melanogaster* homologue of selenophosphate synthetase, is involved in imaginal disc morphogenesis. *Mol. Gen. Genet.*, **257**, 113–123.
- Alsina, B., Corominas, M., Berry, M.J., Bagaña, J. and Serras, F. (1999) Disruption of selenoprotein biosynthesis affects cell proliferation in the imaginal discs and brain of *Drosophila melanogaster*. *J. Cell Sci.*, **112**, 2875–2884.

S. Castellano et al.

- Altschul, S.F., Madden, T., Schaffer, A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*, **25**, 3389–3402.
- Berry, M.J., Harney, J.W., Ohama, T. and Hatfield, D.L. (1994) Selenocysteine insertion or termination: factors affecting UGA codon fate and complementary anticodon:codon mutations. *Nucleic Acids Res.*, **22**, 3753–3759.
- Buettner, C., Harney, J.W. and Berry, M.J. (1999) The *Caenorhabditis elegans* homologue of thioredoxin reductase contains a Selenocysteine Insertion Sequence (SECIS) element that differs from mammalian SECIS elements but directs selenocysteine incorporation. *J. Biol. Chem.*, **274**, 21598–21602.
- Burge, C.B. and Karlin, S. (1998) Finding the genes in genomic DNA. *Curr. Opin. Struct. Biol.*, **8**, 346–354.
- Copeland, P.R., Fletcher, J.E., Carlson, B.A., Hatfield, D.L. and Driscoll, D.M. (2000) A novel RNA binding protein, SBP2, is required for the translation of mammalian selenoprotein mRNAs. *EMBO J.*, **19**, 306–314.
- Flohé, L., Andreesen, J.R., Brigelius-Flohé, R., Maiorino, M. and Ursini, F. (2000) Selenium, the element of the moon, in life on Earth. *Life*, **49**, 411–420.
- Ganther, H.E., (1999) Selenium metabolism, selenoproteins and mechanisms of cancer prevention: complexities with thioredoxin reductase. *Carcinogenesis*, **20**, 1657–1666.
- Gladyshev, V.N., Krause, M., Xu, X.M., Korotkov, K.V., Kryukov, G.V., Sun, Q.A., Lee, B.J., Wooton, J.C. and Hatfield, D.L. (1999) Selenocysteine-containing thioredoxin reductase in *C. elegans*. *Biochem. Biophys. Res. Commun.*, **259**, 244–249.
- Guigó, R., Knudsen, S., Drake, N. and Smith, T.F. (1992) Prediction of gene structure. *J. Mol. Biol.*, **226**, 141–157.
- Hausler, D. (1998) Computational genefinding. *Trends Biochem. Sci.*, *Supplementary Guide to Bioinformatics*, pp. 12–15.
- Hirosawa-Takamori, M., Jäckle, H. and Vorbrüggen, G. (2000) The class 2 selenophosphate synthetase gene of *Drosophila* contains a functional mammalian-type SECIS. *EMBO Rep.*, **1**, 441–446.
- Kryukov, G.V., Kryukov, V.M. and Gladyshev, V.N. (1999) New mammalian selenocysteine-containing proteins identified with an algorithm that searches for Selenocysteine Insertion Sequence Elements. *J. Biol. Chem.*, **274**, 33888–33897.
- Lee, B.J., Rajagopalan, M., Kim, Y.S., You, K.H., Jacobson, K.B. and Hatfield, D. (1990) Selenocysteine tRNA^{[Ser]Sec} gene is ubiquitous within the animal kingdom. *Mol. Cell. Biol.*, **10**, 1940–1949.
- Lehner, C.F. and O'Farrell, P.H. (1990) *Drosophila* cdc2 homologues: a functional homologue is coexpressed with a cognate variant. *EMBO J.*, **9**, 3573–3581.
- Lescure, A., Gautheret, D., Carbon, P. and Krol, A. (1999) Novel selenoproteins identified *in silico* and *in vivo* by using a conserved RNA structural motif. *J. Biol. Chem.*, **274**, 38147–38154.
- Low, S.C. and Berry, M.J. (1996) Knowing when not to stop: selenocysteine incorporation in eukaryote. *Trends Biochem. Sci.*, **21**, 203–208.
- Parra, G., Blanco, E. and Guigó, R. (2000) Geneid in *Drosophila*. *Genome Res.*, **10**, 511–515.
- Persson, B.C., Böck, A., Jäckle, H. and Vorbrüggen, G. (1997) SelD homologue from *Drosophila* lacking selenide-dependant monoselenophosphate synthetase activity. *J. Mol. Biol.*, **274**, 174–180.
- Robinson, D.N. and Cooley, L. (1997) Examination of the function of two kelch proteins generated by stop codon suppression. *Development*, **124**, 1405–1417.
- Stadman, T.C. (1996) Selenocysteine. *Annu. Rev. Biochem.*, **65**, 83–100.
- Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL_W: improving the sensitivity of progressive sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.*, **22**, 4673–4680.
- Tujebajeva, R.M., Ransom, D.G., Harney, J.W. and Berry, M.J. (2000a) Expression and characterization of nonmammalian selenoprotein P in the zebrafish, *Danio rerio*. *Genes Cells*, **5**, 897–903.
- Tujebajeva, R.M., Copeland, P.R., Xu, X-M., Carlson, B.A., Harney, J.W., Driscoll, D.M., Hatfield, D.L. and Berry, M.J. (2000b) Decoding apparatus for eukaryotic selenocysteine insertion. *EMBO Rep.*, **1**, 158–163.
- Ursini, F., Heim, S., Kiess, M., Maiorino, M., Roveri, A., Wissing, J. and Flohé, L. (1999) Dual function of the selenoprotein PHGPx during sperm maturation. *Science*, **285**, 1393–1396.
- Zhou, X., Park, S.I., Moustafa, M.E., Carlson, B.A., Crain, F., Diamond, A.M., Hatfield, D.L. and Lee, B.J. (1999) Selenium metabolism in *Drosophila*. Characterization of the selenocysteine tRNA population. *J. Biol. Chem.*, **274**, 18729–18734.

DOI: 10.1093/embo-reports/kve151